



**For Information Contact:**

Beth Casteel, 703-907-8640

[press@psych.org](mailto:press@psych.org)

Jim Rosack, 703-907-7862

[jrosack@psych.org](mailto:jrosack@psych.org)

**Embargoed Until:**

12:01 am EDT; April 1, 2008

Release No. 08-18

## **Stimulant Treatment for Young Children With ADHD Does Not Lead to Increased Risk of Substance Abuse in Early Adulthood**

*New AJP Study Finds Delayed Treatment, not Early Treatment, is Linked to Later Substance Abuse*

**ARLINGTON, Va. (April 1, 2008)** – Treatment of children with stimulant medications for attention-deficit/hyperactivity disorder (ADHD) does not increase the risk of substance abuse later in adolescence and early adulthood.

A new 17-year study of 176 boys, ages 6 to 12, who were treated with the stimulant medication methylphenidate (marketed as Ritalin and other brand names) shows that the risk of substance abuse during late adolescence and into early adulthood is related to the age at which the boys began taking stimulant medications. For the first time, the new study shows that risk of substance abuse was lowest in the group of boys who began stimulant treatment for ADHD at an earlier age.

Specifically, the rate of drug abuse (excluding alcohol dependence or abuse) in those who had started taking methylphenidate early in the disease process (at age 6 or 7) was similar to the rate in a group of healthy comparison subjects. On the other hand, the rate of drug abuse was statistically significantly higher among those who had begun methylphenidate treatment at later ages, between 8 and 12.

The new findings are detailed in a report which will be published online April 1, 2008, by *The American Journal of Psychiatry (AJP)*, the official journal of the American Psychiatric Association. “Age of Methylphenidate Treatment Initiation in Children With ADHD and Later Substance Abuse: Prospective Follow-Up Into Adulthood” is reported by Salvatore Mannuzza, Ph.D., Rachel G. Klein, Ph.D., Francisco X. Castellanos, M.D., and colleagues at the New York University Child Study Center and the Nathan S. Kline Institute for Psychiatric Research. The report will be available online under *AJP in Advance* (<http://ajp.psychiatryonline.org/pap.dtl>) and will appear in the May 2008 print edition of the journal.

Patients, their families, and the clinical community have all expressed concerns that stimulant treatment in early childhood might lead to later substance use disorders, an idea that is theoretically plausible. Stimulant medications and other drugs of abuse increase concentrations of the neurotransmitter dopamine in an area of the brain called the nucleus accumbens. Research has associated the nucleus accumbens with behavioral reinforcement, including the reinforcing actions of abused drugs.

“Several studies by Barkley, Biederman, Loney, and others have now convincingly demonstrated that stimulant treatment of children with ADHD does not contribute to the later development of drug abuse,” said lead author Salvatore Mannuzza, Ph.D., who is a professor of child and adolescent psychiatry at the New York University School of Medicine. “Our findings suggest that stimulant treatment as early as ages 6 and 7 does not increase the risk for developing substance abuse or antisocial behaviors later in life, and may have long-term beneficial effects.”

Last month, a different report by independent investigators also presented evidence against an association between stimulant treatment and substance abuse. That study, presented online under *AJP in Advance* on March 3, appears along with the new report in the May print edition of *AJP*. The previous study, by Joseph Biederman, M.D., and colleagues at Harvard Medical School and Massachusetts General Hospital, was a 10-year follow-up of boys with ADHD. Biederman and his colleagues followed 42 boys who were treated with stimulants and compared them with 39 who were not. In early adulthood, these two groups showed no significant differences in the rates of alcohol use disorders, nicotine dependence, or other substance-related disorders.

An editorial addressing both reports by Nora Volkow, M.D., director of the National Institute on Drug Abuse, and James Swanson, Ph.D., director of the Child Development Center at the University of California at Irvine, also appears in the May *AJP* print edition. Volkow and Swanson discuss the high risk for substance abuse among individuals with ADHD, pointing out that “[t]he evidence that current clinical practice does not increase later substance use or abuse is comforting. But the failure to document that childhood treatment with stimulant medication is able to decrease the high risk of substance abuse in adulthood is distressing. This highlights the need for the development of integrated treatments that target both ADHD and substance abuse.”

*AJP* Editor-in-Chief Robert Freedman commented, “Early diagnosis and treatment of ADHD continues to be supported as a safe and effective, evidence-backed intervention that helps children while they are in school and later as adults.”

Mannuzza and his colleagues were supported on this study by the National Institute of Mental Health (NIMH) and the National Institute on Drug Abuse (NIDA).

**References:**

Mannuzza S, Klein RG, Truong NL, Moulton JL III, Roizen ER, Howell KH, Castellanos FX: Age of Methylphenidate Treatment Initiation in Children With ADHD and Later Substance Abuse: Prospective Follow-Up Into Adulthood. *Am J Psychiatry* (published online April 1, 2008; doi: 10.1176/appi.ajp.2007.07091465)

Biederman J, Monuteaux MC, Spencer T, Wilens TE, MacPherson HA, Faraone SV: Stimulant Therapy and Risk for Subsequent Substance Use Disorders in Male Adults With ADHD: A Naturalistic Controlled 10-Year Follow-Up Study. *Am J Psychiatry* (published online March 3, 2008; doi: 10.1176/appi.ajp.2007.07091486)

Volkow ND, Swanson JM: Does Childhood Treatment of ADHD With Stimulant Medication Affect Substance Abuse in Adulthood? *Am J Psychiatry* (in press; doi: 10.1176/appi.ajp.2008.08020237)

**Note to Editors:**

If you need more information about the *American Journal of Psychiatry*, including embargoed copies of journal articles, contact Jim Rosack at 703-907-7862 / [jrosack@psych.org](mailto:jrosack@psych.org) or the APA Office of Communications & Public Affairs at 703-907-8640 / [press@psych.org](mailto:press@psych.org). If you would like to receive e-TOCs and other email alerts about *AJP*, visit <http://ajp.psychiatryonline.org/subscriptions/etoc.dtl>.

**About the American Journal of Psychiatry:**

The *American Journal of Psychiatry*, the official journal of the American Psychiatric Association (APA), publishes a monthly print issue with scientific articles submitted by psychiatrists and other scientists worldwide. The editing and peer review processes are conducted independently of any other APA component. Therefore, statements in this press release or the articles in the *Journal*, are not official policy statements of the American Psychiatric Association. The *Journal's* editorial policies conform to the **Uniform Requirements of the International Committee of Medical Journal Editors**, of which it is a member.

*AJP in Advance* is a regular online feature within which original research articles—accepted for publication in *The American Journal of Psychiatry*—are posted ahead of their appearance in print. *AJP in Advance* content has been peer reviewed, copyedited, and approved by authors. Articles in *AJP in Advance* may be cited by using the date they were posted online and their unique digital object identifier (DOI). (See [References](#), above.)

For further information about the Journal visit [www.ajp.psychiatryonline.org](http://www.ajp.psychiatryonline.org).

**About the American Psychiatric Association:**

The American Psychiatric Association is a national medical specialty society whose more than 38,000 physician members specialize in diagnosis, treatment, prevention and research of mental illnesses including substance use disorders. Visit the APA at [www.psych.org](http://www.psych.org) and [www.HealthyMinds.org](http://www.HealthyMinds.org).

###